

Recognizing Generalized Gradient Dynamics by Means of Machine Learning

Vojtěch Votruba, Michal Pavelka

Mathematical Institute, Charles University

June 9, 2025

Introduction

- ▶ Motivation: distilling consistent models from time series
- ▶ Combining generalized gradient dynamics with deep learning
- ▶ Related work:
 - ▶ *Variational Onsager Neural Networks (VONNs): A thermodynamics-based variational learning strategy for non-equilibrium PDEs* — Reina et al, 2022
 - ▶ *Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations* — M. Raissi et al., 2019

Outline

- ▶ Gradient Dynamics
- ▶ Deep Learning
- ▶ Employed design
- ▶ Results:
 - ▶ Overdamped particle
 - ▶ Chemical reactions
 - ▶ Fickian diffusion
- ▶ Future outlook

Generalized Gradient Dynamics

- ▶ Grmela, Öttinger (in the context of GENERIC)
- ▶ Derivation using the Theory of Large Deviations (Mielke, Peletier, Renger, 2015)
- ▶ Let $\mathbf{x} = (x^1, \dots, x^n)$ be n state variables of a purely dissipative system
- ▶ Introduce entropy $S = S(\mathbf{x})$ and the entropic conjugates $\mathbf{x}^* = \frac{\delta S}{\delta \mathbf{x}}$
- ▶ Introduce dissipation potential $\Xi(\mathbf{x}, \mathbf{x}^*)$ such that
 1. $\Xi(\mathbf{x}, \mathbf{x}^*) \geq 0$ and $\Xi(\mathbf{x}, \mathbf{0}) = 0$
 2. $\frac{\delta \Xi}{\delta \mathbf{x}^*} \cdot \mathbf{x}^* \geq 0, \quad \forall \mathbf{x}^*$
 3. near $\mathbf{x}^* = \mathbf{0}$ the potential Ξ must be convex

Generalized Gradient Dynamics

- ▶ Evolution equations

$$\dot{\mathbf{x}} = \left. \frac{\delta \Xi}{\delta \mathbf{x}^*} \right|_{\mathbf{x}^* = \frac{\delta S}{\delta \mathbf{x}}}$$

- ▶ The consequences of imposed restrictions:

- ▶ 2nd law: $\dot{S} = \frac{\delta S}{\delta \mathbf{x}} \cdot \dot{\mathbf{x}} = \mathbf{x}^* \cdot \frac{\delta \Xi}{\delta \mathbf{x}^*} \geq 0$

- ▶ End of evolution at $\mathbf{x}^* = \frac{\delta S}{\delta \mathbf{x}} = \mathbf{0}$

- ▶ \iff Onsager's principle $\frac{\delta S}{\delta \mathbf{x}} = \frac{\delta \Xi^*}{\delta \dot{\mathbf{x}}}$, through Legendre transform

Deep Learning

- ▶ ∈ Machine learning
- ▶ Training and designing deep neural networks
- ▶ Feedforward network: $\mathbf{\Lambda} : \mathbb{R}^N \rightarrow \mathbb{R}^M$

$$\mathbf{\Lambda}(\mathbf{x}; \theta) := \varphi_n(W_n \cdot (\varphi_{n-1}(\dots \varphi_1(W_1 \cdot \mathbf{x} + \mathbf{b}_1) + \mathbf{b}_2)) + \mathbf{b}_n)$$

- ▶ $\theta = \text{weights } \{W_i\}_{i=1}^n + \text{biases } \{b_i\}_{i=1}^n$. φ are *activation functions*
- ▶ Why use neural nets? They are **universal function approximators**

Deep Learning

- ▶ Training := optimizing a loss function $L : \mathbb{R}^M \rightarrow [0, +\infty)$

$$L(\theta) \propto \|\mathbf{y}^{(\text{true})}(\mathbf{x}) - \mathbf{\Lambda}(\mathbf{x}, \theta)\|^2 \implies \frac{\partial L}{\partial \theta} = \mathbf{0}.$$

- ▶ Automatic differentiation engine
- ▶ First-order and Second-order methods
 - ▶ SGD, **Adam**, RMSProp
 - ▶ Newton's method, conjugate gradients, BFGS, **L-BFGS**

Employed Design

- ▶ Variational Onsager Neural Networks
 - ▶ Dissipation neural net – Partially convex
 - ▶ Entropy neural net – Concave (in our case)
 - ▶ Foundation in Amos, Xu, Kolter, 2017
 - ▶ Using reparameterization to fulfill additional restrictions
- ▶ Data generation via numerical integration
- ▶ Calculating $\frac{\delta S}{\delta \mathbf{x}}, \frac{\delta \Xi}{\delta \mathbf{x}^*}$ through automatic differentiation

$$L \propto \|\dot{\mathbf{x}}^{(\text{nn.})} - \dot{\mathbf{x}}^{(\text{true})}\|^2 = \left\| \left. \frac{\partial \Xi^{(\text{nn.})}}{\partial \mathbf{x}^*} \right|_{\mathbf{x}^* = \frac{\partial S^{(\text{nn.})}}{\partial \mathbf{x}}} - \dot{\mathbf{x}}^{(\text{true})} \right\|^2$$

Overdamped Particle


- ▶ Evolution equation $\dot{\mathbf{x}} = -\gamma\mathbf{x}$, where $\gamma > 0$
- ▶ This implies quadratic dissipation potential and entropy

$$S(\mathbf{x}) = -\frac{1}{2}\|\mathbf{x}\|^2, \quad \Xi(\mathbf{x}, \mathbf{x}^*) = \frac{1}{2}\gamma\|\mathbf{x}\|^2$$


- ▶ Evidently, there is non-uniqueness
 - ▶ Addition $S := S + C$, $C \in \mathbb{R}$
 - ▶ Multiplication $S := S/C$ and $\Xi := C\Xi$, where $C > 0$

Overdamped Particle

► Prediction results




(a) 1D




(b) 2D

Overdamped Particle

► Learned functions 1D




(c) Entropy




(d) Dissipation Potential

Overdamped Particle

► Learned functions 2D



(e) Entropy



(f) Dissipation Potential

Chemical Reactions

- ▶ System of N chemical reactions with the concentrations $\mathbf{c} = (c^1, \dots, c^M)$
- ▶ Each reaction looks like



- ▶ Stoichiometric matrix $\nu_{mn} := \beta_m^{(n)} - \alpha_m^{(n)}$
- ▶ Kinetics vector $\Gamma_n = k_f^{(n)} \prod_{m=1}^M c_m^{\alpha_m^{(n)}} - k_b^{(n)} \prod_{m=1}^M c_m^{\beta_m^{(n)}}$
- ▶ Law of mass action: $\dot{\mathbf{c}} = \nu \cdot \Gamma$

Chemical Reactions

▶ Through generalized gradient dynamics:

▶ Entropy: $S(\mathbf{c}) = -\sum_{m=1}^M c_m (\ln c_m + Q_m)$

▶ Dissipation: $\Xi(\mathbf{c}, \mathbf{c}^*) = \sum_{n=1}^N W_n(\mathbf{c}) (\cosh(X_n/2) - 1)$

▶ Where $X_n = -\sum_{m=1}^M c_m^* \nu_{mn}$, $W_n = W_0 \sqrt{\prod_{m=1}^M c_m^{|\nu_{mn}|}}$

▶ Then if we set $k_f^{(n)} := \frac{W_0}{4} \exp\left(-\sum_m \frac{(Q_m+1)\nu_{mn}}{2}\right)$, $k_b^{(n)} := \frac{W_0}{4} \exp\left(\sum_m \frac{(Q_m+1)\nu_{mn}}{2}\right)$

▶ \implies We obtain the right equations

Chemical Reactions


- ▶ Again, non-uniqueness.
- ▶ Example reaction $A \rightleftharpoons B$
 - ▶ According to the law of mass action: $\dot{c}_A = c_B - c_A$, $\dot{c}_B = c_A - c_B$.
 - ▶ This can be clearly satisfied by

$$S(\mathbf{c}) = -\frac{1}{2}\|\mathbf{c}\|^2, \quad \Xi(\mathbf{c}, \mathbf{c}^*) = \frac{1}{2}\mathbf{c}^* \cdot \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \cdot \mathbf{c}^*$$

- ▶ Therefore, fix one quantity to obtain the other


Chemical Reactions

► Example system:




Chemical Reactions

► Prescribed entropy




(g) Evolution for each species



(h) Dissipation potential


Chemical Reactions

► Prescribed dissipation




Chemical Reactions

► Without prescription




Chemical Reactions

► Without prescription



(k) Entropy




(l) Dissipation potential

Fickian Diffusion

- ▶ State vector \rightarrow State function: $c(\mathbf{r})$
- ▶ Linear diffusion equation: $\dot{c}(\mathbf{r}) = \nabla \cdot (D\nabla c(\mathbf{r})) = D\Delta c(\mathbf{r})$
- ▶ Through generalized gradient dynamics:
 - ▶ Entropy $S[c(\mathbf{r})] = -\int c \log(c) d^3r$
 - ▶ Dissipation $\Xi[c(\mathbf{r}), \nabla c^*(\mathbf{r})] = \frac{1}{2} \int Dc \|\nabla c^*\|^2 d^3r$
- ▶ 1D discretization $\Delta c \approx \frac{c_{i+1} - 2c_i + c_{i-1}}{h^2}$, where h is the grid resolution
- ▶ Training on Fourier modes with PBC, evaluation on 3 different initial conditions


Fickian Diffusion

► Example training trajectory




Fickian Diffusion

► Polynomial initial conditions




Fickian Diffusion

► Discontinuous initial conditions



Fickian Diffusion

► Gaussian spikes initial conditions



Future Outlook

- ▶ Integrating with reversible networks, i.e., Deep Poisson Neural Networks
 - ▶ Building an entire GENERIC coupling $\dot{\mathbf{x}} = \{\mathbf{x}, H\} + \frac{\delta \Xi}{\delta \mathbf{x}^*} \Big|_{\mathbf{x}^* = \frac{\delta S}{\delta \mathbf{x}}}$
- ▶ Extending to functional spaces via Neural Operators (Kovachki, Li et al, 2023)
- ▶ Very open to collaboration!
 - ▶ Vojtěch Votruba, votruba@karlin.mff.cuni.cz
 - ▶ Michal Pavelka, pavelka@karlin.mff.cuni.cz

Thank you for your attention.